

European clinical guidelines for Tourette Syndrome and other tic disorders. Part III: behavioural and psychosocial interventions

Cara Verdellen · Jolande van de Griendt ·
Andreas Hartmann · Tara Murphy ·
the ESSTS Guidelines Group

© Springer-Verlag 2011

Abstract This clinical guideline provides recommendations for the behavioural and psychosocial interventions (BPI) of children and adolescents with tic disorders prepared by a working group of the European Society for the Study of Tourette Syndrome (ESSTS). A systematic literature search was conducted to obtain an update on the efficacy of BPI for tics. Relevant studies were identified using computerised searches of the Medline and PsycINFO databases and the Cochrane Library for the years 1950–2010. The search identified no meta-analyses, yet twelve (systematic) reviews and eight randomised controlled trials provided evidence for the current review. Most evidence was found for habit reversal training (HRT) and the available but smaller evidence also supports the efficacy of exposure with response prevention (ERP). Both interventions are considered first line behavioural treatments for tics for both children and adults and should be offered to a patient, taking into account his preference. Treatments that are considered second line or add-on behavioural treatments are contingency management, function based interventions and relaxation training. Neurofeedback is still experimental.

Members of the ESSTS Guidelines Group are listed under [Appendix](#).

C. Verdellen (✉) · J. van de Griendt
HSK Group/Expertise Centre Tics, Hambakenwetering 5B,
5231 DD Den Bosch, The Netherlands
e-mail: c.verdellen@hsk.nl

A. Hartmann
Centre de Référence National ‘Syndrome Gilles de la Tourette’,
Pôle des Maladies du Système Nerveux, Groupe Hospitalier
Pitié-Salpêtrière, Assistance Publique des Hôpitaux de Paris,
Paris, France

T. Murphy
Tourette Syndrome Clinic, Great Ormond Street Hospital NHS
Trust, London, UK

Almost no research was identified that examined the efficacy of psychosocial interventions, e.g., psychoeducation and group work. Based on clinical practice, this guideline recommends behavioural treatment as first line offer to patients in most cases. It should be embedded within a psychoeducational and supportive context and can be combined with drug treatment.

Keywords Tourette · Tic disorders · Guidelines · Behavioural treatment · Psychosocial interventions · Literature review

Introduction

Tic disorders (e.g., chronic motor or vocal tic disorder and Tourette syndrome; TS) are a common developmental disorder among children and adolescents. About 10% of children have (motor and/or vocal) tics during childhood, which in many cases disappear. However, approximately 1% of children and adolescents develop TS [1], a condition that is characterised by multiple motor and vocal tics. Often, symptoms of Attention Deficit Hyperactivity Disorder (ADHD), Obsessive Compulsive Disorder (OCD) and Pervasive Developmental Disorder (PDD) coexist. Tics may vary in their severity and in the inconvenience they cause to an individual and his environment. Although not all children and families seek help [2], many require some form of treatment or support to manage their tics and impaired social, emotional and behavioural functioning. The European Society for the Study of Tourette Syndrome (ESSTS) met in November 2008 and in August 2009 to discuss the development of guidelines for the treatment of tic disorders. The present guideline describes recommendations for the behavioural and psychosocial interventions

(BPI) of tics in children and adolescents. The behavioural treatment of comorbidity in tic disorders is beyond the scope of this guideline, but some information can be found in Döpfner and Rothenberger [3, 4] and Rothenberger et al. [5].

Method

To obtain an update on scientific research on the efficacy and effectiveness of BPI for tic disorders, a literature search was performed on behavioural and psychosocial interventions using computerised searches of the Medline (PubMed) and PsycINFO databases (1950–May 2010). This search was performed using relevant terms (e.g., tics, tic disorder, Tourette's syndrome, behavio(u)r therapy, habit reversal, exposure and response prevention, relaxation, psychoeducation, support group, coping strategies). 'Map term to subject heading', in which the best-matching subject headings associated with a term are included, was used. In addition, the reference lists of the (review) articles identified through Medline and PsychINFO were reviewed for additional studies. The search produced an initial list of 159 publications, which revealed no meta-analyses, twelve (systematic) reviews, eight randomised controlled trials (RCT's) and one non-randomised controlled trial. Furthermore, multiple case studies were found. No studies were found on tic disorders comparing behaviour therapy with pharmacotherapy. The Cochrane library was also consulted and contained no information on BPI for tic disorders. Although the publication list comprises many relevant articles related to the subject, it should be noted that it is not exhaustive.

Results

Behavioural interventions

The most rigorously researched of all psychological interventions are behavioural treatments. Overall, eight different behavioural treatments for tics were identified in the literature: massed (negative) practice (MP), habit reversal (HR), self-monitoring (SM), contingency management (CM)/function based interventions (FBI), relaxation training (RT)/hypnosis, exposure and response prevention (ER), cognitive behavioural treatment (CBT) and bio(neuro) feedback (NF). Most studies utilised single-case experimental designs, using a multiple baseline design, a waiting list design (WL) or an alternating treatments design. Nine studies employed group-based methodology, eight of these with random assignment to treatment conditions. RCT's differed from one another on several variables (e.g., tic disorder, HR ingredients, sample age, outcome measures,

duration of follow-up). Four studies included both children and adults, but only two studies consisted of children (age 7–18) only. Table 1 provides an overview of the RCT's of tic disorders.

Massed (negative) practice (MP)

MP is the oldest behavioural treatment for tics reported in the literature. MP involves the repeated, rapid, voluntary and effortful performance of tics for a specified period of time (e.g., 30 min), interspersed with brief periods of rest. The majority of single-case studies into MP failed to find a decrease in tic frequency. In the one RCT comparing MP to HR in both children and adults with TS, MP failed to produce reductions in tics comparable to those produced by the HR condition [6] (see Table 1). Overall, it can be concluded that MP has limited therapeutic value for tic disorders.

Habit Reversal (HR)

HR is the most extensively researched behavioural treatment for tics. It offers a set of techniques that help the patient become aware of the occurrence of a tic, followed by a so-called competing response training to interrupt or inhibit the tic [7–9]. HR may also comprise relaxation training, contingency management, and generalization training. Awareness of the tics is enhanced through response description, response detection, an early warning procedure, and situation awareness training. With competing response training the patient learns to initiate a response for 1–3 min, or until the urge goes away, contingent upon the urge to perform a tic or after the actual occurrence of the tic. Several dozens of uncontrolled studies into HR report tic reductions of 30–100%. Most studies involved small patient samples with unspecified diagnostic criteria. However, eight RCT's (see Table 1) were performed into HR indicating its' efficacy. Four studies were performed on adults only (age 18–60) [10–13], 1 study (the largest study up to date, N = 126) included children only (age 9–17) [14] and 3 studies included both children and adults (age 6–62) [6, 8, 15]. Most studies consisted of 10–14 weekly sessions [10–13, 15]. Some studies spreaded the sessions over time after starting with weekly sessions, e.g. biweekly or monthly sessions. Follow-up data of at least 10 months duration showed maintenance of results. More information can be found in Table 1. In addition to the RCT's, several review articles support evidence for HR [16–21]. Awareness training and competing response training seem to be the active components of the multi-component HR package that are necessary and sufficient for tic reduction [22]. In summary, studies indicate that HR is effective for both

Table 1 Randomised controlled trials of behavioural interventions for tics disorders (divided according to the inclusion or exclusion of children)

Study	Interventions	Subjects	Outcome	Efficacy	Follow-up	Comments
Azrin et al. [6]	HR, MP parallel	22 TS patients (age 11–62, average 30.5)	HR > MP**	Tic frequency (self-report of number of tic episodes): HR: 84% decrease MP: ±30% decrease	1 month: HR: 92% decrease MP: ±30% decrease 4–6 months: HR: 99% decrease 18 months: HR: 97% decrease No follow-up	<ul style="list-style-type: none"> • 1 or 2 sessions (2.5 h) • HR consisted of competing response training and general behaviour therapy procedures • FU: telephone calls • No further FU data on MP
Azrin and Peterson [8]	HR, WL Crossover	10 TS patients (age 6–36, average 18.1)	HR > WL*	Tic frequency (self-report and independent video rating): HR: 93% decrease at home (self-report); 93.5% at clinic (independent video rating) WL: no significant decrease	No follow-up	<ul style="list-style-type: none"> • Mean of 20 sessions (range 13–30) during 8–11 months • HR consisted of awareness training, self-monitoring, competing response training, relaxation training and contingency management
Verdellen et al. [15]	HR, ER parallel	43 TS patients (age 7–55, average 20.6)	HR = ER	YGTSS: ER: 26.2 pre 17.6 post (ES 1.42) 58% of patients showed >30% reduction HR: 24.1 pre 19.7 post (ES 1.06) 28% of patients showed >30% reduction Tic frequency clinic (independent video rating): ER: 74% of patients showed >30% reduction (ES 0.90) HR: 53% of patients showed >30% reduction (ES 0.47)	3 months: YGTSS: ER: 14.0 (ES 1.49) HR: 13.5 (ES 1.95)	<ul style="list-style-type: none"> • 12 sessions ER (2 h, weekly) vs 10 sessions HR (1 h, weekly) • Weighted gain scores were used to correct for number of session • HR consisted of awareness training and competing response training • HR and ER improved significantly on all measurements • FU included 25 patients who had received the alternative condition
Bergin et al. [29]	RT, MT parallel	23 TS patients (age 7–18, average 11.3)	RT = MT	Tic frequency home (report of parent/partner): ER: 89% of patients showed >30% reduction (ES 0.88) HR: 72% of patients showed >30% reduction (ES 0.73) YGTSS: RT: 86% of patients improved (non-significant) MT: 66% of patients improved (non-significant)	6 weeks: YGTSS: RT: 71% of patients improved (non-significant) MT: 55% of patients improved (non-significant)	<ul style="list-style-type: none"> • 6 sessions (1 h, weekly) • No significant improvements • Comparable results on other tic severity measurements (Goetz Videotape Scale, Hopkins Motor and Vocal Tic Scale, Tourette Syndrome Severity Scale)

Table 1 continued

Study	Interventions	Subjects	Outcome	Efficacy	Follow-up	Comments
Piacentini et al. [14]	HR, SP parallel	126 TS/CTD patients (age 9–17, average 11.7)	HR > SP	YGTSS: HR: 24.7 pre- 17.1 post (ES 0.68) SP: 24.6 pre- 21.1 post	3 months: YGTSS: HR: 13.9 (n = 23) SP: 9.9 (n = 6) 6 months: YGTSS: HR: 13.3 (n = 23) SP: 10.4 (n = 6)	<ul style="list-style-type: none"> • 8 sessions (during 10 weeks; first two sessions 1.5 h, remaining sessions 1 h) • 3 monthly booster sessions for responders • HR consisted of awareness training, competing response training, relaxation training and function based interventions • SP consisted of supportive psychotherapy and education. • Significantly more children receiving HR were rated as being very much improved or much improved on the Clinical Global Impression—improvement scale (52.5 vs 18.5%) • 3 and 6 months follow-up: results remained stable for the available responders
Wilhelm et al. [10]	HR, SP parallel	32 TS patients (average age 36.2)	HR > SP*	YGTSS: HR: 30.5 pre- 19.8 post (ES 1.50) SP: 26.6 pre- 26.9 post (ES -0.03)	10 months: YGTSS: HR: 21.0 (still significantly improved) SP: 23.8 (pre-treatment level)	<ul style="list-style-type: none"> • 14 sessions (8 weekly, 6 2-weekly) • HR consisted of self-monitoring, competing response training, relaxation training and contingency management • HR improved significantly on the Clinical Global Impression—improvement scale (CGI-i) as well, SP not
Deckersbach et al. [11]	HR, SP parallel	30 TS patients (average age 35.1)	HR > SP*	YGTSS: HR: 39.3 pre- 18.3 post SP: 27.7 pre- 26.8 post	6 months: YGTSS: HR: 18.4 SP: 26.6	<ul style="list-style-type: none"> • 10 months FU: no significant difference between HR and SP on the YGTSS; significant difference between HR and SP maintained on the CGI-i • 14 sessions (8 weekly, 4 2-weekly, 2 monthly) • HR consisted of awareness training, self-monitoring, competing response training, relaxation training and contingency management • Both groups improved in life-satisfaction and psychosocial functioning; only HR improved significantly in tic severity • 6 months FU: results remained stable

Table 1 continued

Study	Interventions	Subjects	Outcome	Efficacy	Follow-up	Comments
O'Connor et al. [12]	HR, CBT Parallel	14 CTD patients (age 23–49)	HR = CBT	Tic frequency (diary): HR: 54% decrease CBT: 57% decrease Degree of control: HR: 65% CBT: 78%	3 months: Tic frequency (diary): HR: 77% decrease CBT: 86% decrease Degree of control: HR: 69% CBT: 87% 2 years: Degree of control: HR: 43% of patients had maintained degree of control CBT: 50% of patients had maintained degree of control	<ul style="list-style-type: none"> • 10 sessions (weekly) • HR consisted of awareness training, competing response training and contingency management • CBT included HR elements; cognitive approach showed no significant additional effect • 2 years FU: interview by telephone
O'Connor et al. [13]	HR/CBT, WL Crossover	47 CTD patients (age 18–60, average 39.1)	HR/CBT > WL*	Tic frequency and intensity (diary, video): HR/CBT: 88% of patients significant decrease in frequency and intensity on diary and video measures WL: no significant decrease Degree of control: HR/CBT: 65% of patients reported between 75 and 100% control	2 months: Tic frequency, intensity and degree of control: HR/CBT: maintained results 2 years: Degree of control: HR/CBT: 52% of patients maintained 75–100% control	<ul style="list-style-type: none"> • Group non-RCT design • 12 sessions (weekly) • Both tic disorders (n = 47) and habit disorders (n = 43) were included • HR consisted of awareness training, competing response training, relaxation training, relapse prevention and was accompanied with more general cognitive and behavioural restructuring • 2 years FU: interview by telephone

Significance of unbolditalics—children & Adults, bold—children only, italics—adults only

TS Tourette Syndrome, CTD chronic tic disorder, MP massed practice, WL waiting list, HR habit reversal, ER exposure and response prevention, RT relaxation training, MT minimal therapy, SP supportive psychotherapy, CBT cognitive behavioural therapy, YGTSS Yale Global Tic Severity Scale, ES effect size, FU follow-up

* Significant difference $p < 0.05$, ** significant difference $p < 0.01$

vocal and motor tics, for children as well as adults, for patients receiving TS medications as well as those not doing so, for tic severity as well as tic frequency, and with no evidence of symptom substitution [23].

Self-monitoring (SM)

SM consists of having the patient record tics during a specified period by using a counter or a notebook. The goal is to identify when and in which situation a tic occurs. SM is often applied in multi-component treatment (HR) packages. Only a few case studies have been conducted in which SM was used as the primary treatment procedure, indicating that SM generally results in temporary improvement [24]. Immediate, rather than delayed recording of tics appears to be essential for tic reduction [25, 26]. In summary, there is some support for the usefulness of SM, although the generalizability and durability of treatment gains remain unclear.

Contingency management (CM)/Function based interventions (FBI)

CM consists of the manipulation of environmental contingencies, so that tic-free intervals are positively reinforced and tic behaviours are ignored. Using tokens to reinforce the absence of tics is an example of a CM-based intervention [27]. CM is mostly used in multi-component treatment packages, making it difficult to assess the unique value of this specific technique. A more specialised type of CM is known as ‘function based interventions’ (FBI). In FBI, factors specific to the individual’s unique environment are identified which increase or decrease tics. The factors are altered to provide tic reductions. The effect of this intervention is reported in single-case studies [28].

Relaxation training (RT)

The use of RT in the treatment of tics is based on the observation that increases in stress and anxiety result in concomitant increases in tic performance. Hence, tic reduction in RT might work indirectly via stress reduction. Relaxation can help reduce muscular tension. RT includes deep breathing, progressive muscle training, and imagery, and is mostly applied as part of a multi-component treatment. Two studies examined RT as a mono-component treatment under controlled conditions. Peterson and Azrin [24] found that RT resulted in an average tic decrease of 32% ($N = 6$), which was less than the effect of competing response training (55% decrease) or SM (45% decrease). Bergin et al. [29] conducted a controlled, randomised trial in children (age 7–18) comparing RT with minimal control therapy. After 6 weekly sessions, they found no differences

between groups in terms of tic symptom severity (see Table 1). Improvements in tic symptoms were only short term. In summary, it can be concluded that RT is not an evidence-based treatment for tics, although it may have some merit in the short term. Hypnosis and cue-controlled relaxation, more specific forms of relaxation, have been shown useful in some cases [30, 31] and need further study.

Exposure and response prevention (ER)

The application of ER to reduce tics is based on the association of unpleasant premonitory sensations followed by a motor or vocal tic that relieves the sensation [32]. In learning theory terms, tics can be viewed as conditioned responses to premonitory, interoceptive stimuli. When such stimuli reoccur over time, the power of the associative interaction between the sensations and the resultant tic behaviour is strengthened. ER aims at interrupting the association, thus preventing the tics to occur. By confronting patients for a prolonged period of time with the sensations (exposure) and resisting the tic (response prevention), the patients might learn to tolerate the unpleasant sensation (habituation). Habituation will lessen the urge or need to give into the tic, resulting in a reduction of tic behaviour. One RCT comparing 12 2-h ER sessions (including 2 training sessions) with 10 1-h HR sessions [15] revealed no differences between conditions (see Table 1). The study included both children and adults (age 7–55) with TS. Effect sizes were larger for ER than for HR. Results of this RCT and of a few case studies into ER [33] provide support for ER and indicate the need for further study. In addition, review articles [17–20] suggest that ER is a promising treatment for tics. Preliminary results indicate that ER is effective for vocal and motor tics, for children as well as adults, for tic severity as well as tic frequency, with no indications of a rebound effect [34]. Since younger children are less aware of premonitory sensory motor phenomena it has to be clarified if there is an age effect respective an age limit for ER but also for HR [35].

Cognitive behavioural treatment (CBT)

Two studies (both included adults only) were identified that employed cognitive behavioural interventions (see Table 1). The cognitive component of the intervention consisted of mentally challenging and restructuring the way patients evaluated their expectations and actions in high-risk situations related to frequent ‘ticking’ behaviour. One study compared CBT with HR in CTD and found comparable results for both treatments [12]. In another study, CBT combined with HR in CTD was found superior to WL [13]. The value of cognitive interventions in the

treatment of tics remains to be investigated separately, as studies into CBT included a HR component, thus making its unique contribution to the results unclear. Preliminary results seem to indicate that cognitive interventions have no specific additive value in the treatment of tics.

Bio(neuro) feedback (NF)

NF is aimed at self-directed modulating of defined parts of brain electrical activity [36]. It is based on operant conditioning; when brain activity changes in the desired direction, a positive ‘reward’ feedback is given to the individual.

Two case studies on TS (both $N = 2$) were found, one with positive results [37], the other with inconclusive results [38]. In addition, a study in 15 TS patients showed significantly lower tics during relaxation biofeedback compared to arousal biofeedback [39]. Considering the lack of randomised controlled research on NF and biofeedback for the treatment of tics, further research is required in order to draw conclusions.

Psychosocial interventions

The literature search and recent reviews by Cook and Blacher [17] and Woods et al. [40] indicate there is no evidence-based practice with regard to supportive psychosocial interventions in TS. There is limited literature guiding interventions which modify the child’s environment (parents, teachers, peers) to facilitate reduction of tics and enhance quality of life [41, 42]. However, education about tic disorders and the provision of support and reassurance to patients and their families have been described as the cornerstone for all other treatment interventions [43]. Therefore, based on clinical practice rather than empirical evidence, the present guideline includes recommendations on these approaches.

Psychoeducation

It is broadly accepted that TS and related tic disorders affect the lives of patients and all family members. Specifically, family members often have difficulty adjusting to the diagnosis and may struggle to manage with comorbid difficulties such as anger outbursts and hyperactivity [44]. Psychoeducation represents an important component in providing support and resolving misunderstanding, uncertainty and stigma in TS [45]. Information about the natural course of TS and helping the child and his environment to identify personal strengths often serves to reduce anxiety. It is proposed to provide relevant information to a child about the condition. This will promote coping strategies and self-efficacy and provide the child with the tools to explain to others (especially teachers and schoolmates) about the

disorder. Providing educators with general information relating to the aetiology, presentation, and course of TS will help them to implement effective individualised strategies to manage classroom behaviour, thereby allowing the child to maximise learning potential. Due to the rapidly changing nature of TS, Kepley and Connors [46] highlight the need for a flexible approach, close liaison between parents and teachers and frequent re-evaluation of interventions within education to ensure effective management.

Children with TS report that they struggle to fit into society’s expectations and rate associated embarrassment as disabling as the tics themselves [45, 47]. Young people with TS have been rated as less socially favourably as other non-affected peers in studies using laboratory-based paradigms [48–50]. Two studies have looked at peer-perceptions of TS and the effects of psychoeducation through a videotape intervention [49, 51]. In one study, older students viewed a TS-specific educational video. These students reported more positive attitudes toward people with tics than those who viewed other non-TS related material [51]. Another study involving younger children showed no difference in attitudes following presentation of psychoeducational material. The effect on social behaviour intentions and actual social behaviour remains unclear and further research in this area is needed.

Group work

Case studies have been published describing and evaluating group work targeting different symptoms in children with TS. Group approaches offer children and their parents the opportunity to meet one another within a supportive environment. Groups can provide psychoeducational material [52] on aspects of TS such as tic management, anger management, bullying and school and self-esteem. Although the approach remains at the descriptive stage, qualitative feedback following such interventions has been positive and it seems worthy of further evaluation.

Charities and support organisations

The preponderance of charities for people with TS throughout the world are a vehicle for sharing information on TS. Taubert [53] discusses the important role of voluntary organisations in supporting people with TS and their families. Large organisations such as the Tourette Syndrome Association in the US provide care and advocacy. Meanwhile, several European self-help organisations (e.g. in Germany, France, the Netherlands) are also quite active. While support groups for families are important, a more holistic approach between professionals, charities and government bodies may also be key role for such organisations.

Conclusions

The literature review provides scientific evidence for the behavioural treatment of tics in tic disorders. Most studies investigated the effectiveness of HR. Based on eight RCTs, several review articles and dozens of case studies, HR can be considered an evidence-based behavioural treatment for tics for both children and adults. HR was found superior to WL [8, 13], MP [6], and SP [10, 11, 14], and equally effective as ER [15]. A number of RCTs have shown that the improvement of HR continues up to at least 10 months following completion of treatment. The literature also provides support for ER. In one RCT [15] comparing ER with HR, larger effect sizes were found for ER than for HR, indicating its efficacy. In addition, several review articles and a few case studies demonstrate that ER is a promising, probably efficacious treatment for tic disorders for both children and adults, which needs further research.

Although most evidence was found for HR and, to a lesser extent, for ER, the current literature base is limited by the availability of only one (albeit large-scale) RCT [14] with a child and adolescent-only sample. Therefore, firm conclusions for these specific age groups cannot be drawn. Comparability of the RCT's was limited by the different outcome measures that were used, application of different HR ingredients, different sample ages, number of sessions, and follow-up periods. Only three RCT's [10, 14, 15] reported effect sizes. In order to enable comparisons of efficacy, future studies are encouraged to include such calculations. There are no controlled studies into the efficacy and effectiveness of other psychosocial interventions, e.g., psychoeducation or support groups. Therefore, these interventions are not to be considered evidence based, although the importance of these interventions is emphasised in clinical practice.

Recommendations


Based on the evidence at present and clinical support this guideline recommends both HR and ER as first line behavioural treatments for tics for both children and adults. HR has a broader evidence base and is usually offered. Results indicate that ER may be more effective than HR at the severe end of TS, i.e., when many tics are involved [15]. However, ER as applied in the RCT took twice as much therapy time as HR, i.e., results were based on sessions lasting 2 h each, whereas HR sessions usually last 1 h. Recent research shows that ER sessions lasting for 1 h had comparable results [54]. Other treatments for tics were identified in the literature, for which insufficient evidence was found. MP, the oldest behavioural treatment for tics reported in the literature, lacks evidence and therefore is not recommended as a treatment for tics. CM/

reinforcement techniques and FBI are also not to be considered evidence-based treatments for tics. However, case studies indicate that these interventions have some value in specific cases and need further research. CM may be a useful tool for children, because motivation for a behavioural treatment may not be intrinsic and its reward (i.e., tic reductions) may be too far away for a child. Based on the current literature, CM and FBI are recommended as second line behavioural interventions. RT, although also not to be considered an evidence-based treatment for tics, is recommended as a second line behavioural treatment for tics as well. Case studies indicate that patients can benefit from this treatment to some extent and hypnosis and cue-controlled relaxation were found to reduce tics via stress reduction in specific cases. Cognitive interventions, although these have not been studied in adolescents, do not seem to have additive value to HR. Therefore, at present, this method is not recommended as an intervention for tics. Based on the sparse and inconclusive literature on the efficacy of NF, this method cannot be recommended as a treatment for tics. However, a RCT in children with ADHD showed a long-term improvement on self-regulation of behaviour, which seems promising also for TS, especially in cases where TS and ADHD may coexist [55, 56]. NF should thus be further evaluated as a treatment for tics.

Although there are no controlled studies into the efficacy and effectiveness of other psychosocial interventions, e.g., psychoeducation, bibliotherapy, self-help, social support groups, this guideline recommends embedding each treatment within a psychoeducational and supportive context. Psychoeducation alone may also be useful for families who do not engage with more comprehensive treatments due to very mild tics or lack of services. Specific interventions must target not only tic symptoms, but also comorbid problems and coping strategies that can profoundly influence the impact that TS may have on an individual's well being during childhood and later into adulthood. The present guideline does not provide recommendations as to the behavioural treatment of comorbidity in tic disorders.

A summary of the recommendations of this guideline is presented in a flowchart (see Fig. 1). Psychoeducation is recommended in all (both mild and more severe) cases. Other supportive interventions (e.g., liaison with school, group work, social skills training) are recommended in individual cases depending on the specific needs of a child and his environment. HR and ER are behavioural treatments recommended as first line treatments to patients who want to reduce their tics. In helping a child or adolescent (and his family) making a choice which treatment to start with, information should be given about both treatments. This guideline recommends to evaluate treatment response after 10 sessions, and to continue treatment for another 10 sessions when there is a partial response, to start the

Fig. 1 Decision tree of psychological interventions for tics in tic disorders



alternative treatment when there is no response, and to consider medication when both treatments have not resulted in relevant tic reductions. In case of severe symptoms, it is recommended to consider combining a behavioural treatment with medication or starting medication first and adding behavioural treatment later. If the combined treatment is effective it is advised to consider phasing out the medication. In addition, second line interventions such as CM, FBI or RT are recommended in individual cases, when first line interventions do not suffice.

Conflict of interest Commercial firms and governmental organisations did not play a role in, or fund, the development of these guidelines. Cara Verdellen, Jolande van de Griendt, Andreas Hartmann, Tara Murphy declare that they have no conflict of interest.

Appendix: Members of the ESSTS Guidelines Group

Christos Androutsos, Harald Aschauer, Gillian Baird, Netty Bos-Veneman, Ariana Brambilla, Francesco Cardona, Danielle C. Cath, Andrea E. Cavanna, Virginie Czernecki, Sandra Dehning, Alan Eapter, Luca Farkas, Julia Gadaros, Andreas Hartmann, Elizabeth Hauser, Isabel Heyman, Tammy Hedderly, Pieter J. Hoekstra, Anne Korsgaard, Georgina M. Jackson, Linnea Larsson, Andrea G. Ludolph, Davide Martino, Claudia Menghetti, Nanette Mol Debes, Norbert Muller, Kirsten Muller-Vahl, Alexander Munchau,

Tara Murphy, Richard Musil, Peter Nagy, Judith Nurnberger, Ben Oostra, Perry Paschou, Massimo Pasquini, Kirsten J. Plessen, Mauro Porta, Hugh Rickards, Renata Rizzo, Mary M. Robertson, Veit Roessner, Aribert Rothenberger, Domenico Servello, Liselotte Skov, Jeremy S. Stern, Gerd Strand, Zsannett Tarnok, Cristiano Termine, Jolande Van der Griendt, Cara Verdellen, Veerle Visser-Vandewalle, Ebba Wannag, Tomas Wolanczyk

References

- Robertson MM (2008) The prevalence and epidemiology of Gilles de la Tourette syndrome. Part 1: the epidemiological and prevalence studies. *J Psychosom Res* 65(5):461–472. doi: [10.1016/j.jpsychores.2008.03.006](https://doi.org/10.1016/j.jpsychores.2008.03.006)
- Schlender M, Schwarz O, Rothenberger A, Roessner V (2010) Tic disorders: administrative prevalence and co-occurrence with attention-deficit/hyperactivity disorder in a German community sample. *Eur Psychiatry*. doi: [10.1016/j.eurpsy.2009.10.003](https://doi.org/10.1016/j.eurpsy.2009.10.003)
- Döpfner M, Rothenberger A (2007) Behavior therapy in tic-disorders with co-existing ADHD. *Eur Child Adolesc Psychiatry* 16(Suppl 1):89–99. doi: [10.1007/s00787-007-1011-7](https://doi.org/10.1007/s00787-007-1011-7)
- Döpfner M, Rothenberger A (2007) Tic- und Zwangsstörungen; Themenschwerpunkt. *Kindheit und Entwicklung* 16:75–138
- Rothenberger A, Roessner V, Banaschewski T (2006) Habit formation in Tourette Syndrome with associated obsessive-compulsive behaviour: at the crossroads of neurobiological modelling. *Behav Brain Sci* 29:627–628
- Azrin NH, Nunn RG, Frantz SE (1980) Habit reversal vs negative practice treatment of nervous tics. *Behav Ther* 11(2):169–178

7. Azrin NH, Nunn RG (1973) Habit reversal: a method of eliminating nervous habits and tics. *Behav Res Ther* 11:619–628
8. Azrin NH, Peterson AL (1990) Treatment of Tourette Syndrome by habit reversal: a waiting-list control group comparison. *Behav Ther* 21(3):305–318
9. Woods DW, Miltenberger RG (1995) Habit reversal: a review of applications and variations. *J Behav Ther Exp Psychiatry* 26(2): 123–131
10. Wilhelm S, Deckersbach T, Coffey BJ, Bohne A, Peterson AL, Baer L (2003) Habit reversal versus supportive psychotherapy for Tourette's disorder: a randomized controlled trial. *Am J Psychiatry* 160(6):1175–1177
11. Deckersbach T, Rauch S, Buhlmann U, Wilhelm S (2006) Habit reversal versus supportive psychotherapy in Tourette's disorder: a randomized controlled trial and predictors of treatment response. *Behav Res Ther* 44(8):1079–1090. doi:10.1016/j.brat.2005.08.007
12. O'Connor K, Gareau D, Borgeat F (1997) A comparison of a behavioural and a cognitive-behavioural approach to the management of chronic tic disorders. *Clin Psychol Psychother* 4:105–117
13. O'Connor KP, Brault M, Robillard S, Loiselle J, Borgeat F, Stip E (2001) Evaluation of a cognitive-behavioural program for the management of chronic tic and habit disorders. *Behav Res Ther* 39(6):667–681
14. Piacentini J, Woods DW, Scahill L, Wilhelm S, Peterson AL, Chang S, Ginsburg GS, Deckersbach T, Dziura J, Levi-Pearl S, Walkup JT (2010) Behavior therapy for children with Tourette disorder: a randomized controlled trial. *JAMA* 303(19):1929–1937. doi:10.1001/jama.2010.607
15. Verdellen CW, Keijsers GP, Cath DC, Hoogduin CA (2004) Exposure with response prevention versus habit reversal in Tourette's syndrome: a controlled study. *Behav Res Ther* 42(5):501–511. doi:10.1016/S0005-7967(03)00154-2
16. Carr JE, Chong IM (2005) Habit reversal treatment of tic disorders: a methodological critique of the literature. *Behav Modif* 29(6):858–875. doi:10.1177/0145445505279238
17. Cook CR, Blacher J (2007) Evidence-based psychosocial treatments for tic disorders. *Clin Psychol Sci Pract* 14(3):252–267
18. Himle MB, Woods DW, Piacentini JC, Walkup JT (2006) Brief review of habit reversal training for Tourette syndrome. *J Child Neurol* 21(8):719–725
19. Peterson AL (2007) Psychosocial management of tics and intentional repetitive behaviors associated with Tourette syndrome. In: Woods DW, Piacentini JC, Walkup JT (eds) *Treating Tourette syndrome and tic disorders: a guide for practitioners*. Guilford Press, New York, pp 154–184
20. Piacentini J, Chang S (2001) Behavioral treatments for Tourette syndrome and tic disorders: state of the art. *Adv Neurol* 85:319–331
21. Piacentini J, Chang S (2005) Habit reversal training for tic disorders in children and adolescents. *Behav Modif* 29(6):803–822. doi:10.1177/0145445505279385
22. Miltenberger RG, Fuqua RW, McKinley T (1985) Habit reversal with muscle tics: replication and component analysis. *Behav Ther* 16(1):39–50
23. Woods DW, Conelea CA, Walther MR (2007) Barriers to dissemination: exploring the criticisms of behavior therapy for tics. *Clin Psychol Sci Pract* 14:279–282
24. Peterson AL, Azrin NH (1992) An evaluation of behavioral treatments for Tourette syndrome. *Behav Res Ther* 30(2):167–174
25. Billings A (1978) Self-monitoring in the treatment of tics: a single-subject analysis. *J Behav Ther Exp Psychiatry* 9(4):339–342
26. Turpin G, Powell GE (1984) Effects of massed practice and cue-controlled relaxation on tic frequency in Gilles de la Tourette's syndrome. *Behav Res Ther* 22(2):165–178
27. Woods DW, Himle MB (2004) Creating tic suppression: comparing the effects of verbal instruction to differential reinforcement. *J Appl Behav Anal* 37(3):417–420. doi:10.1901/jaba.2004.37-417
28. Watson TS, Sterling HE (1998) Brief functional analysis and treatment of a vocal tic. *J Appl Behav Anal* 31(3):471–474. doi: 10.1901/jaba.1998.31-471
29. Bergin A, Waranch HR, Brown J, Carson K, Singer HS (1998) Relaxation therapy in Tourette syndrome: a pilot study. *Pediatr Neurol* 18(2):136–142
30. Culbertson FM (1989) A four-step hypnotherapy model for Gilles de la Tourette's syndrome. *Am J Clin Hypn* 31:252–256
31. Hoogduin K, Cazemier B, Verdellen C (1995) Exposure, responspreventie en zelfcontrole bij Gilles de la Tourettepatiënten [Exposure, response prevention and self-control in patients with Gilles de la Tourette syndrome]. *Gedragstherapie* 28:31–40
32. Leckman JF, Walker DE, Cohen DJ (1993) Premonitory urges in Tourette's syndrome. *Am J Psychiatry* 150(1):98–102
33. Hoogduin C, Verdellen C, Cath D (1997) Exposure and response prevention in the treatment of Gilles de la Tourette's syndrome: four case studies. *Clin Psychol Psychother* 4:125–137
34. Verdellen CW, Hoogduin CA, Keijsers GP (2007) Tic suppression in the treatment of Tourette's syndrome with exposure therapy: the rebound phenomenon reconsidered. *Mov Disord* 22(11):1601–1606. doi:10.1002/mds.21577
35. Banaschewski T, Woerner W, Rothenberger A (2003) Premonitory sensory phenomena and suppressibility of tics in Tourette Syndrome—developmental aspects in children. *Dev Med Child Neurol* 45:700–703
36. Heinrich H, Gevensleben H, Strehl U (2007) Annotation: neurofeedback—train your brain to train behaviour. *J Child Psychol Psychiatry* 48(1):3–16. doi:10.1111/j.1469-7610.2006.01665.x
37. Tansey MA (1986) A simple and a complex tic (Gilles de la Tourette's syndrome): their response to EEG sensorimotor rhythm biofeedback training. *Int J Psychophysiol* 4(2):91–97
38. Breteler MHM, van Heeswijk L, Arns M, Verdellen C (2008) Psychologische interventies bij het syndroom van Gilles de la Tourette [Psychological interventions in Gilles de la Tourette syndrome]. *Tijdschrift voor Neuropsychiatrie en Gedragsneurologie* 7:158–164
39. Nagaia Y, Cavanna A, Critchley HD (2009) Influence of sympathetic autonomic arousal on tics: Implications for a therapeutic behavioral intervention for Tourette syndrome. *J Psychosom Res* 67:599–605
40. Woods DW, Piacentini JC, Walkup JT (2007) *Treating Tourette syndrome and tic disorders: a guide for practitioners*. Guilford Press, New York
41. Conelea CA, Woods DW (2008) The influence of contextual factors on tic expression in Tourette's syndrome: a review. *J Psychosom Res* 65(5):487–496. doi:10.1016/j.jpsychores.2008.04.010
42. Packer LE (2005) Tic-related school problems: impact on functioning, accommodations, and interventions. *Behav Modif* 29(6):876–899. doi:10.1177/0145445505279383
43. Peterson BS, Cohen DJ (1998) The treatment of Tourette's syndrome: multimodal, developmental intervention. *J Clin Psychiatry* 59(Suppl 1):62–72 discussion 73–64
44. Dooley JM, Berna PM, Gordon KE (1999) Parent perception of symptom severity in Tourette's syndrome. *Arch Dis Child* 81:440–441
45. Cutler D, Murphy T, Gilmour J, Heyman I (2009) The quality of life of young people with Tourette syndrome. *Child Care Health Dev* 35(4):496–504. doi:10.1111/j.1365-2214.2009.00983.x
46. Kepley HO, Connors S (2007) Management of learning and school difficulties in children with Tourette syndrome. In: Woods DW, Piacentini JC, Walkup JT (eds) *Treating Tourette syndrome*

- and tic disorders: a guide for practitioners. Guilford Press, New York, pp 242–264
47. Wand RR, Matazow GS, Shady GA, Furer P, Staley D (1993) Tourette syndrome: associated symptoms and most disabling features. *Neurosci Biobehav Rev* 17(3):271–275
 48. Boudjouk PJ, Woods DW, Miltenberger RG, Long ES (2000) Negative peer evaluation in adolescents: effects of tic disorders and trichotillomania. *Child Fam Behav Ther* 22:17–28
 49. Friedrich S, Morgan SB, Devine C (1996) Children's attitudes and behavioral intentions toward a peer with Tourette syndrome. *J Pediatr Psychol* 21(3):307–319
 50. Woods DW, Fuqua RW, Outman RC (1999) Evaluating the social acceptability of persons with habit disorders: the effects of topography, frequency, and gender manipulation. *J Psychopathol Behav Assess* 21:1–18
 51. Woods DW, Marcks BA (2005) Controlled evaluation of an educational intervention used to modify peer attitudes and behavior toward persons with Tourette's Syndrome. *Behav Modif* 29(6):900–912. doi:[10.1177/0145445505279379](https://doi.org/10.1177/0145445505279379)
 52. Murphy T, Heyman I (2007) Group work in young people with Tourette Syndrome. *Child Adolesc Mental Health* 12:46–48
 53. Taubert KA (1999) Role of voluntary organizations in clinical care research and public policy. In: Leckman JF, Cohen DJ (eds) *Tourette's syndrome-tics obsessions compulsions: developmental psychopathology and clinical care*. Wiley, Hoboken, pp 399–413
 54. Griendt Jvd, Verdellen C, Verbraak M, Hoogduin C (Manuscript in preparation) The effect of exposure length on treatment outcome for tics in Tourette's syndrome: One hour sessions versus two hour sessions
 55. Gevensleben H, Holl B, Albrecht B, Vogel C, Schlamp D, Kratz O, Studer P, Rothenberger A, Moll GH, Heinrich H (2009) Is neurofeedback an efficacious treatment for ADHD? A randomised controlled clinical trial. *J Child Psychol Psychiatry* 50(7):780–789. doi:[10.1111/j.1469-7610.2008.02033.x](https://doi.org/10.1111/j.1469-7610.2008.02033.x)
 56. Gevensleben H, Holl B, Albrecht B, Schlamp D, Kratz O, Studer P, Rothenberger A, Moll GH, Heinrich H (2010) Neurofeedback training in children with ADHD: 6-month follow-up of a randomised controlled trial. *Eur Child Adolesc Psychiatry* 19(9):715–724. doi:[10.1007/s00787-010-0109-5](https://doi.org/10.1007/s00787-010-0109-5)